arXiv:1303.4327v1 [math.AG] 18 Mar 2013

Homogeneous division polynomials for Weierstrass elliptic curves

Jinbi Jin
Universiteit Leiden
Mathematisch Instituut
Niels Bohrweg 1
2333 CA Leiden
jjin@math.leidenuniv.nl

March 20, 2013

0 Introduction

Abstract. Starting from the classical division polynomials we construct homogeneous polynomials α_n, β_n, γ_n such that for $P = (x : y : z)$ on an elliptic curve in Weierstrass form over an arbitrary ring we have $nP = \big(\alpha_n(P) : \beta_n(P) : \gamma_n(P)\big)$. To show that $\alpha_n, \beta_n, \gamma_n$ indeed have this property we use the a priori existence of such polynomials, which we deduce from the Theorem of the Cube.

Acknowledgements. This work is partially funded by NWO, the Netherlands Organisation for Scientific Research. This work was originally a part of my MSc thesis, and I would like to thank Bas Edixhoven for his supervision at the time, especially for mentioning the Theorem of the Cube. I would also like to thank both Bas Edixhoven and Lenny Taelman for their many comments on previous (unpublished) versions of this paper.

Introduction. The main problem treated in this paper is the following.

> *Given an integer n, construct a triple of homogeneous polynomials that defines multiplication by n on all* Weierstrass curves *(i.e. projective plane curves given by a Weierstrass polynomial) over rings.*

The classical *division polynomials* describe multiples of *affine points* — these are points of the form $(a : b : 1)$ — for all smooth Weierstrass curves over fields. This is a basic result that is stated in several books on elliptic curves. An example of a book that treats the theory behind this in more detail is [3]. This result is not completely satisfactory, as the division polynomials are not defined on non-affine points.

In this paper we construct for all integers n three homogeneous polynomials α_n, β_n, γ_n of degree n^2 such that for all Weierstrass curves C over a ring R, and all “nowhere singular” points on C of the form $P = (x : y : z)$, we have $nP = \big(\alpha_n(P) : \beta_n(P) : \gamma_n(P)\big)$. See the example on the next page for α_2, β_2, γ_2.

Overview. In Section 1.1, we define Weierstrass curves over arbitrary schemes, and give some properties of them that we will use throughout this paper. In Section 1.2, we describe the construction of α_n, β_n, γ_n. In Section 1.3, we state the main theorem. Section 1.2 and 1.3, as well as the summary of Section 1.1, should be accessible for anyone with some basic knowledge about elliptic curves over fields and commutative algebra.

In Section 2.1, we consider multiplication by n on smooth Weierstrass curves, by reducing to the *universal smooth Weierstrass curve* and use the Theorem of the Cube to show the existence of three homogeneous polynomials of degree n^2 defining multiplication by n on smooth Weierstrass curves. We compare these polynomials with α_n, β_n, γ_n, and show that they are the same. In Section 2.2, we show that α_n, β_n, γ_n in fact define multiplication by n on all Weierstrass curves, using the fact that they do so on the universal smooth Weierstrass curve.

Example 0.1 (Doubling formula). Let R be a ring, and let C be the Weierstrass curve over R defined by the polynomial

$$W = y^2z + a_1xyz + a_3yz^2 - x^3 - a_2x^2z - a_4xz^2 - a_6z^3,$$

where $a_1, a_2, a_3, a_4, a_6 \in R$. Let $P = (x : y : z)$ be a point on C given in homogeneous coordinates, such that $(\partial W/\partial x)(P)$, $(\partial W/\partial y)(P)$, $(\partial W/\partial z)(P)$ generate the unit ideal in R. Then

$$2P = \big(\alpha_2(P) : \beta_2(P) : \gamma_2(P)\big),$$

where

$$\begin{aligned}
\alpha_2 = {} & 2xy^3 + 3a_1x^2y^2 + (a_1^2 - 2a_2)y^3z + (a_1^3 - 3a_1a_2 + 3a_3)xy^2z + (-2a_1^2a_2 + 2a_2^2 - 6a_4)x^2yz \\
& + (a_1a_2^2 - 3a_2a_3 - 3a_1a_4)y^2z^2 + (a_1^2a_2^2 - a_1^3a_3 - 2a_1a_2a_3 - 4a_1^2a_4 - 3a_3^2 + 2a_2a_4 - 18a_6)xyz^2 \\
& + (-a_1a_2^3 + a_1^2a_2a_3 + a_2^2a_3 - 3a_1a_3^2 + 4a_1a_2a_4 - 3a_3a_4 - 9a_1a_6)x^2z^2 \\
& + (a_1a_2^2a_3 - a_1^2a_3^2 - 3a_2a_3^2 - a_1a_3a_4 - 3a_1^2a_6 + 2a_4^2 - 6a_2a_6)yz^3 \\
& + (-a_1a_2a_3^2 - a_1a_2^2a_4 + 2a_1^2a_3a_4 - a_1^3a_6 - 2a_3^3 + a_2a_3a_4 + 4a_1a_4^2 - 3a_1a_2a_6 - 9a_3a_6)xz^3 \\
& + (-a_2a_3^3 + a_1a_3^2a_4 - a_1a_2^2a_6 + a_3a_4^2 - 3a_2a_3a_6 + 3a_1a_4a_6)z^4, \\
\beta_2 = {} & y^4 + a_1xy^3 + (a_1a_2 - 2a_3)y^3z + (a_1^2a_2 - a_2^2 - 3a_1a_3 + 3a_4)xy^2z \\
& + (-2a_1a_2^2 + 6a_1a_4)x^2yz + (a_2^3 - a_1a_2a_3 + a_1^2a_4 - 5a_2a_4 + 18a_6)y^2z^2 \\
& + (a_1a_2^3 - 2a_1^2a_2a_3 + a_1^3a_4 - a_2^2a_3 + 3a_1a_3^2 - 6a_1a_2a_4 + 3a_3a_4 + 27a_1a_6)xyz^2 \\
& + (-a_2^4 + 2a_1a_2^2a_3 - a_1^2a_2a_4 + 6a_2^2a_4 - 6a_1a_3a_4 + 9a_1^2a_6 - 9a_4^2)x^2z^2 \\
& + (a_2^3a_3 - a_1a_2a_3^2 + a_1^3a_6 + 2a_3^3 - 5a_2a_3a_4 - a_1a_4^2 + 3a_1a_2a_6 + 18a_3a_6)yz^3 \\
& + (a_1^2a_2a_3^2 - a_1^3a_3a_4 + a_1^4a_6 + 2a_2^2a_3^2 - a_1a_3^3 - a_2^3a_4 - 2a_1^2a_4^2 + 6a_1^2a_2a_6 - 6a_3^2a_4 + 3a_2a_4^2 + 9a_2^2a_6 - 27a_4a_6)xz^3 \\
& + (a_1a_2a_3^3 - a_1^2a_3^2a_4 + a_1^3a_3a_6 - a_3^4 + a_2a_3^2a_4 - 2a_1a_3a_4^2 - a_2^3a_6 + 6a_1a_2a_3a_6 - a_4^3 - 9a_3^2a_6 + 9a_2a_4a_6 - 27a_6^2)z^4, \\
\gamma_2 = {} & 8y^3z + 12a_1xy^2z + 6a_1^2x^2yz + (a_1^3 + 12a_3)y^2z^2 + (a_1^4 + 12a_1a_3)xyz^2 + (-a_1^3a_2 + 3a_1^2a_3)x^2z^2 \\
& + (a_1^3a_3 + 6a_3^2)yz^3 + (-a_1^3a_4 + 3a_1a_3^2)xz^3 + (-a_1^3a_6 + a_3^3)z^4.
\end{aligned}$$

1 Definitions, statement of main theorem

Rings in this paper are always assumed to be commutative, associative, and unital.

1 Weierstrass curves over schemes

For now, we will use the language of schemes. For a summary of this section in terms of Weierstrass curves over rings, we refer to the end of this section.

Let S be a scheme. A *Weierstrass curve* C over S is a closed subscheme of $\mathbb{P}^2_S$ defined by a homogeneous polynomial of the form

$$W = y^2z + a_1xyz + a_3yz^2 - x^3 - a_2x^2z - a_4xz^2 - a_6z^3 \tag{1}$$

with $a_1, a_2, a_3, a_4, a_6 \in \mathcal{O}_S(S)$. The *discriminant* $\Delta \in \mathcal{O}_S(S)$ of C is the discriminant of the equation (1), see e.g. [3, Def. 2.7].

We now describe the functor of points of the Weierstrass curve C explicitly. If T is an S-scheme, $\mathcal{L}$ is an invertible $\mathcal{O}_T$-module, $s_0, s_1, s_2 \in \mathcal{L}(T)$, and $f \in \mathcal{O}_S(S)[x,y,z]$ homogeneous of degree d, then we have a well-defined section $f(s_0, s_1, s_2) \in \mathcal{L}^{\otimes d}(T)$. We say that s_0, s_1, s_2 *generate* $\mathcal{L}$ if $(s_0)_x, (s_1)_x, (s_2)_x$ generate $\mathcal{L}_x$ as an $\mathcal{O}_{T,x}$-module for all $x \in T$. The following is a special case of a well-known result, see e.g. [7, Lem. 01O4], as the graded $\mathcal{O}_S$-algebra $\mathcal{O}_S[x,y,z]/(W)$ is generated by its degree 1 part.

Proposition 1.1. *Let S be a scheme, and let C be a Weierstrass curve over S defined by W as in (1). Then for all S-schemes T, we have*

$$C(T) = \left\{ (\mathcal{L}, s_0, s_1, s_2) : \begin{array}{c} \mathcal{L} \text{ invertible } \mathcal{O}_T\text{-module, } s_0, s_1, s_2 \in \mathcal{L}(T) \text{ generating } \mathcal{L} \\ W(s_0, s_1, s_2) = 0 \text{ in } \mathcal{L}^{\otimes 3} \end{array} \right\} \Big/ \cong .$$

Here, $(\mathcal{L}, s_0, s_1, s_2) \cong (\mathcal{M}, t_0, t_1, t_2)$ if and only if there exists an isomorphism $\mathcal{L} \to \mathcal{M}$ of $\mathcal{O}_T$-modules mapping s_i to t_i for all $i \in \{0, 1, 2\}$.

If $\mathcal{L} = \mathcal{O}_T$, then we denote the class of $(\mathcal{L}, s_0, s_1, s_2)$ by $(s_0 : s_1 : s_2)$.

Proposition 1.2. *Let S be a scheme, and let C be a Weierstrass curve over S defined by W as in (1). Then C is flat, proper, and locally of finite presentation over S, and its geometric fibres are integral curves of arithmetic genus 1.*

Proof. Except for flatness, our claim is well-known. We check affine locally on the base that C is flat over S. So assume $S = \operatorname{Spec} R$. Then C is covered by the two standard affine open covers in which y and z, respectively, are invertible. Since W is monic with respect to x, the corresponding R-algebras are free as R-modules, hence flat. We deduce that C is flat over S. ☐

We deduce that C is smooth (of relative dimension 1) if and only if C is smooth on all geometric fibres, i.e. $\Delta \neq 0$ in all geometric points of S (see e.g. [3, Prop. 2.25]). This holds if and only if $\Delta \in \mathcal{O}_S(S)^\times$.

In general, let C^{sm} be the smooth locus of C over S. Then we have the following description of the functor of points of C^{sm}.

Proposition 1.3. *Let S be a scheme, and let C be a Weierstrass curve over S defined by W as in (1). Then for all S-schemes T, the set $C^{\mathrm{sm}}(T)$ is the subset of $C(T)$ consisting of the classes of the 4-tuples $(\mathcal{L}, s_0, s_1, s_2)$ such that the sections*

$$(\partial W/\partial x)(s_0, s_1, s_2), (\partial W/\partial y)(s_0, s_1, s_2), (\partial W/\partial x)(s_0, s_1, s_2) \in \mathcal{L}^{\otimes 2}(T)$$

generate $\mathcal{L}^{\otimes 2}$.

Proof. We denote the structure morphism of C by f. Note that for all $x \in C$, we have that C is smooth in x if and only if $C_{f(x)}$ is smooth in x, using Proposition 1.2. The polynomials $\partial W/\partial x, \partial W/\partial y, \partial W/\partial z$ define the singular locus on every fibre, so C^{sm} is the open subscheme of C that is the complement of the common zero locus of these polynomials. $\square$

By [2, Prop. II.2.7], we have the following.

Theorem 1.4. *Let S be a scheme, and let C be a smooth Weierstrass curve over S. Then there exists a unique commutative group scheme structure on C with zero section $(0:1:0) \in C(S)$.*

In general, note that $(0:1:0) \in C^{\mathrm{sm}}(S)$, since $(\partial W/\partial z)(0,1,0) = 1$ generates $\mathcal{O}_S$. To show that every Weierstrass curve C over a scheme S admits a commutative group scheme structure on C^{sm} with zero section $(0:1:0) \in C^{\mathrm{sm}}(S)$, we first consider the *universal Weierstrass curve.*

Let $\mathcal{A} = \mathbb{Z}[a_1, a_2, a_3, a_4, a_6]$. Then the *universal Weierstrass curve $\mathcal{C}$* is the Weierstrass curve over $\mathcal{A}$ defined by

$$\mathcal{W} = y^2z + a_1xyz + a_3yz^2 - x^3 - a_2x^2z - a_4xz^2 - a_6z^3 \in \mathcal{A}[x, y, z].$$

It is universal in the following sense.

Proposition 1.5. *Let S be a scheme, and let C be a Weierstrass curve over S defined by W as in (1). Then there exist unique morphisms $g\colon C \to \mathcal{C}$ and $h\colon S \to \operatorname{Spec}\mathcal{A}$ such the following diagram is commutative, and such that the outer square is Cartesian.*

$$\begin{tikzcd}
C \arrow[r] \arrow[d, "g"'] & \mathbb{P}^2_S \arrow[r] \arrow[d, dashed, "h_{\mathbb{P}^2_{\mathcal{A}}}" description] & S \arrow[d, "h"] \\
\mathcal{C} \arrow[r] & \mathbb{P}^2_{\mathcal{A}} \arrow[r] & \operatorname{Spec}\mathcal{A}
\end{tikzcd}$$

Proof. It suffices to show this in the case that S is affine; the general case follows from it by gluing.

So suppose that $S = \operatorname{Spec} A$, and let ϕ denote the morphism $S \to \operatorname{Spec}\mathcal{A}$ given by the ring morphism $\mathcal{A} \to A$ sending every a_i to the corresponding coefficient of W. Then $C = \mathcal{C} \times_{\mathcal{A}} S$ (via ϕ), so $(\phi_{\mathcal{C}}, \phi)$ is a pair satisfying the required properties.

Suppose that (g, h) is a pair of morphisms satisfying the required properties. Then note that $\mathcal{C} \to \mathbb{P}^2_{\mathcal{A}}$ is a closed immersion, hence a monomorphism, so it follows that g is the unique morphism making the diagram commute. Moreover, $h_{\mathbb{P}^2_{\mathcal{A}}}$ induces a map from C to $\mathcal{C}$ if and only if the induced map $\mathcal{O}_{\mathbb{P}^2_{\mathcal{A}}}(3)(\mathbb{P}^2_{\mathcal{A}}) \to \mathcal{O}_{\mathbb{P}^2_{A}}(3)(\mathbb{P}^2_{A})$ (which is the map $\mathcal{A}[x,y,z]_3 \to A[x,y,z]_3$ induced by the ring morphism $\mathcal{A} \to A$) sends $\mathcal{W}$ to W, in other words, if and only if $h = \phi$. $\square$

We now have the following corollary of Theorem 1.4.

Corollary 1.6. *There exists a unique commutative group scheme structure on $\mathcal{C}^{\mathrm{sm}}$ with zero section $(0:1:0) \in \mathcal{C}^{\mathrm{sm}}(S)$.*

Proof. First note that by Proposition 1.2 and [2, Prop. II.2.7], $\mathcal{C}^{\mathrm{sm}}$ admits a group scheme structure with zero section $(0:1:0) \in \mathcal{C}^{\mathrm{sm}}(\mathcal{A})$. We show that it must be unique.

Note that $\mathcal{C}^{\mathrm{sm}} \times_{\mathcal{A}} \mathcal{C}^{\mathrm{sm}}$ is smooth over $\mathcal{A}$, with irreducible fibres, and that $\mathcal{A}$ is integral. Hence (for example by combining [7, Lem. 004Z] and [4, Prop. 17.5.7]) $\mathcal{C}^{\mathrm{sm}} \times_{\mathcal{A}} \mathcal{C}^{\mathrm{sm}}$ is an integral scheme. Also note that $\mathcal{C}^{\mathrm{sm}}$ is separated over $\mathcal{A}$.

Now consider $\mathcal{R} = \mathbb{Z}[a_1, a_2, a_3, a_4, a_6, 1/\Delta]$. Then $\operatorname{Spec} R$ is an open subscheme of $\operatorname{Spec} A$. Let $\mathcal{E} = \mathcal{C} \times_{\mathcal{A}} \operatorname{Spec} \mathcal{R}$, and note that it is a non-empty open subscheme of $\mathcal{C}^{\mathrm{sm}}$. As $\mathcal{E}$ is a smooth Weierstrass curve, $\mathcal{E}$ admits a unique group scheme structure with zero section $(0:1:0) \in \mathcal{E}(\mathcal{R})$. Hence the group scheme structure on $\mathcal{C}^{\mathrm{sm}}$ extends that of $\mathcal{E}$, so by the above this must be the unique extension of the group scheme structure of $\mathcal{E}$ to $\mathcal{C}^{\mathrm{sm}}$. ☐

By universality of $\mathcal{C}$, this gives a natural commutative group scheme structure on every Weierstrass curve.

We now summarise in terms of Weierstrass curves over rings.

Summary. Let R be a ring. A *Weierstrass curve* C over R is a closed subscheme of $\mathbb{P}^2_R$ defined by a homogeneous polynomial of the form (1) with $a_1, a_2, a_3, a_4, a_6 \in R$. The *discriminant* $\Delta \in R$ of C is the discriminant of the equation (1), see e.g. [3, Def. 2.7]. The Weierstrass curve C is smooth if and only if $\Delta \in R^\times$.

We have an explicit description of the set $C(R)$ of R-valued points of C as follows. If M is an invertible R-module, $m_0, m_1, m_2 \in M$, and $f \in R[x,y,z]$ homogeneous of degree d, then we have $f(s_0, s_1, s_2) \in M^{\otimes d}$. Then

$$C(R) = \left\{ (M, m_0, m_1, m_2) : \begin{array}{c} M \text{ invertible } R\text{-module, } m_0, m_1, m_2 \in M \\ Rm_0 + Rm_1 + Rm_2 = M,\ W(m_0, m_1, m_2) = 0 \text{ in } M^{\otimes 3} \end{array} \right\} \Big/ \cong,$$

where $(M, m_0, m_1, m_2) \cong (N, n_0, n_1, n_2)$ if and only if there exists an isomorphism $M \to N$ of R-modules mapping m_i to n_i for all $i \in \{1, 2, 3\}$. If $M = R$, the class of (M, m_0, m_1, m_2) is denoted $(m_0 : m_1 : m_2)$.

As an example, if R is a field (or more generally, any ring with trivial Picard group), then

$$C(R) = \{(x,y,z) \in R^3 : W(x,y,z) = 0\}/R^\times.$$

In general, let C^{sm} be the open subscheme of C that is the complement of the common zero locus of $\partial W/\partial x, \partial W/\partial y, \partial W/\partial x$. Then $C^{\mathrm{sm}}(R)$ is the subset of $C(R)$ consisting of the classes of the 4-tuples (M, m_0, m_1, m_2) such that

$$R(\partial W/\partial x)(m_0, m_1, m_2) + R(\partial W/\partial y)(m_0, m_1, m_2) + R(\partial W/\partial z)(m_0, m_1, m_2) = M^{\otimes 2}.$$

The set $C^{\mathrm{sm}}(R)$ contains the point $(0:1:0) \in C(R)$, and has a natural structure of an abelian group with $(0:1:0)$ as neutral element.

2 Division polynomials and construction of α_n, β_n, γ_n

Let us recall the definition of division polynomials. The main reference for this is [3, Ch. 3]. For this purpose, we consider a special (but yet also generic!) smooth Weierstrass curve.

Let K be the algebraic closure of the field of fractions of the ring $\mathcal{A} = \mathbb{Z}[a_1, a_2, a_3, a_4, a_6]$. Let E be the elliptic curve over K defined by the homogeneous Weierstrass polynomial

$$\mathcal{W} = y^2 z + a_1 xyz + a_3 yz^2 - x^3 - a_2 xz^2 - a_4 x^2 z - a_6 z^3.$$

Further, let X, Y denote the rational functions x/z and y/z, respectively. Then the function field $K(E)$ of E is the field of fractions of $K[X,Y]/(\mathcal{W}')$, where

$$\mathcal{W}' = Y^2 + a_1 XY + a_3 Y - X^3 - a_2 X^2 - a_4 X - a_6$$

is the affine Weierstrass polynomial. We have a leading coefficient map $\Lambda\colon K(E)-0 \to K-0$, given by $f \mapsto \big((X/Y)^{-\mathrm{ord}_0 f} f\big)(0)$, where $\mathrm{ord}_0 f$ denotes the order of f at the point at infinity of E. This is well-defined as X/Y has a simple zero at 0.

Then we can define the division polynomials as follows.

Definition 1.7. Let $n \in \mathbb{Z}-\{0\}$. The *n-th division polynomial* Ψ_n is the unique rational function f in $K(E)$ with divisor $\sum_{P\in E[n]-0}\langle P\rangle - (n^2-1)\langle 0\rangle$ and leading coefficient n. Moreover, we define Ψ_0 to be 0.

These division polynomials exist, as the degree of the divisors defining them is 0, and for all $n \in \mathbb{Z}-0$, we have $\sum_{P\in E[n]} P = 0$.

Remark 1.8. The division polynomials satisfy the following relation for all $m, n \in \mathbb{Z}$ (see [3, Prop. 3.53])

$$\Psi_{m+n}\Psi_{m-n} = \Psi_{m+1}\Psi_{m-1}\Psi_n^2 - \Psi_{n+1}\Psi_{n-1}\Psi_m^2.$$

Using Ψ_1, Ψ_2, Ψ_3, and Ψ_4, and the recurrence relation above, one can recursively compute the other division polynomials.

Furthermore, define the following rational functions.

$$\Phi_n = X\Psi_n^2 - \Psi_{n-1}\Psi_{n+1} \qquad \Omega_n = \begin{cases} 1 & \text{if } n=0 \\ \frac{1}{2\Psi_n}\big(\Psi_{2n} - \Psi_n^2(a_1\Phi_n + a_3\Psi_n^2)\big) & \text{otherwise} \end{cases}$$

We list some facts in the next proposition.

Proposition 1.9 ([3, Sect. 3.6], [1, Prop. 1.7.7])**.** *Let $n \in \mathbb{Z}$. Then the following are true.*

- $\Psi_n, \Phi_n, \Omega_n \in \mathcal{A}[X,Y]/(\mathcal{W}')$
- $\Lambda\Phi_n = \Lambda\Omega_n = 1$
- $\mathrm{ord}_0\Phi_n = -2n^2$, $\mathrm{ord}_0\Omega_n = -3n^2$

Now we can state the basic result mentioned in the introduction more precisely, in the case of our special smooth Weierstrass curve E/K.

Proposition 1.10 ([3, Prop. 3.55])**.** *Let $P = (x : y : 1) \in E(K)$ be a rational point, and let $n \in \mathbb{Z}$. Then*

$$nP = \big(\Psi_n(x,y)\Phi_n(x,y) : \Omega_n(x,y) : \Psi_n^3(x,y)\big).$$

We want to obtain a version of this result that works for all rational points, including the point at infinity. We will do this by choosing representatives in $\mathcal{A}[X,Y]$ of the elements $\Psi_n\Phi_n$, Ω_n, Ψ_n^3 of $\mathcal{A}[X,Y]/(\mathcal{W}')$ and then homogenising them.

However, one has to be careful with the choice of representatives. As an example of this, the usual representatives chosen for the Ψ_n have Y-degree at most 1, which is convenient for calculations. But this also causes homogenisations to be divisible by high powers of z for large n, as the leading term with respect to y in the homogeneous Weierstrass equation $\mathcal{W}$ is y^2z. This causes problems at the point at infinity for $|n| \geq 2$.

Note that $\mathcal{W}$ is monic with respect to x, which motivates the following definition. Recall that $X = x/z$ and $Y = y/z$.

Definition 1.11. Let $A_n, B_n, C_n \in \mathcal{A}[X,Y]$ be the unique representatives of $\Psi_n\Phi_n$, Ω_n, Ψ_n^3 with X-degree at most 2, and define the polynomials $\alpha_n = A_n z^{n^2}$, $\beta_n = B_n z^{n^2}$, $\gamma_n = C_n z^{n^2}$ in $\mathcal{A}[x,y,z,z^{-1}]$.

Uniqueness of A_n, B_n, C_n is guaranteed by the fact that $\mathcal{W}'$ is monic in X.

Proposition 1.12. *Let $n \in \mathbb{Z}$. Then $\alpha_n, \beta_n, \gamma_n \in \mathcal{A}[x,y,z]$ and are homogeneous of degree n^2. Moreover, $\alpha_n, \gamma_n \in (x,z)$ and $\beta_n \in y^{n^2} + (x,z)$.*

Proof. To show that $\alpha_n, \beta_n, \gamma_n \in \mathcal{A}[x,y,z]$, it suffices to show that the polynomials A_n, B_n, C_n have total degree at most n^2. Note that the monomials with X-degree at most 2 all have distinct orders at infinity. As A_n, B_n, C_n have orders $-3n^2+1$, $-3n^2$, $-3n^2+3$ at infinity, respectively, and as they have X-degree at most 2, it follows that they have total degree at most n^2. (Here, we use that X and Y have poles of orders 2 and 3 at the point at infinity, respectively.)

Now $\alpha_n, \beta_n, \gamma_n$ are by construction homogeneous of degree n^2, and as Y^{n^2} is the unique monomial with order $-3n^2$, the result follows. □

For our special smooth Weierstrass curve E, the problem at the point at infinity is now solved.

Proposition 1.13. *Let $n \in \mathbb{Z}$, and let $P = (x : y : z) \in E(K)$. Then*

$$nP = \big(\alpha_n(P) : \beta_n(P) : \gamma_n(P)\big).$$

Proof. As α_n, β_n, γ_n were obtained from $\Psi_n\Phi_n$, Ω_n, Ψ_n^3 by homogenisation, the result follows from Proposition 1.10 for P of the form $(x : y : 1)$. For $P = (0 : 1 : 0)$, the claim follows from Proposition 1.12. □

3 Statement of the main theorem

Let R be a ring. Given W in $R[x,y,z]$ of the form (1), there is an obvious ring morphism $\mathcal{A} \to R$ which maps $a_i \in \mathcal{A}$ to the corresponding coefficient of W in R. We will view R as an $\mathcal{A}$-algebra via this morphism.

Theorem 1.14. *Let R be a ring, and let C be a Weierstrass curve over R defined by W as in (1). Let $P = [(M, m_0, m_1, m_2)] \in C^{\mathrm{sm}}(R)$. Then*

$$nP = \Big[\big(M^{\otimes n^2}, \alpha_n(m_0,m_1,m_2), \beta_n(m_0,m_1,m_2), \gamma_n(m_0,m_1,m_2)\big)\Big]$$

In Section 2, we will state and prove the scheme-theoretic equivalent of this theorem.

If $M = R$, then $M^{\otimes n^2} = R$, so if a point P is of the form $(x : y : z)$, then nP is also of this form, namely,

$$nP = \big(\alpha_n(x,y,z) : \beta_n(x,y,z) : \gamma_n(x,y,z)\big)$$

However, even for a smooth Weierstrass curve C, the subset of R-valued points of the form $(x : y : z)$ need not be closed under addition, as we will see in Example 1.16.

As α_n, β_n, γ_n were obtained from $\Phi_n\Psi_n$, Ω_n, Ψ_n^3 by homogenisation, we then have the following immediate consequence of Theorem 1.14.

Corollary 1.15. *Let $n \in \mathbb{Z}$, and let C be a Weierstrass curve over a ring R. Then for all $P \in C^{\mathrm{sm}}(R)$ of the form $(x : y : 1)$, we have*

$$nP = \big(\Phi_n(x,y)\Psi_n(x,y) : \Omega_n(x,y) : \Psi_n^3(x,y)\big).$$

Example 1.16. We give an example of a Weierstrass curve C over a ring R and two R-valued points $P, Q \in C(R)$ of the form $(x : y : z)$ such that its sum $P + Q \in C(R)$ is not of the form $(x : y : z)$. Let $R = \mathbb{Z}\big[\sqrt{-5}\big]$, let $K = \mathbb{Q}\big(\sqrt{-5}\big)$, and consider the smooth Weierstrass curve C over R given by

$$W = y^2z + xyz + yz^2 - x^3 - 4xz^2 + 6z^3,$$

and the two points

$$P = (9 : 23 : 1) \qquad\qquad Q = \big(3411\sqrt{-5} : 26488 + 117\sqrt{-5} : -3645\sqrt{-5}\big)$$

in $C(K)$. Note that P and Q define points of the form $(x : y : z)$ in $C(R)$ as well, as the R-submodule of K generated by their coordinates are trivial, but that the R-submodule of K generated by the coordinates of

$$P + Q = \big(61028487 + 104922279\sqrt{-5} : 120011054 - 171672039\sqrt{-5} : -127263527\big) \in C(K)$$

is invertible, but not trivial, so $P + Q$ is not of the form $(x : y : z)$ in $C(R)$.

2

Proof of the main theorem

Here is the scheme-theoretic equivalent of Theorem 1.14.

Theorem 2.1. *Let S be a scheme, and let C be a Weierstrass curve over S defined by W as in (1). Let T be an S-scheme, and let $P = [(\mathcal{L}, s_0, s_1, s_2)] \in C^{\mathrm{sm}}(T)$. Then*

$$nP = \left[\left(\mathcal{L}^{\otimes n^2}, \alpha_n(s_0, s_1, s_2), \beta_n(s_0, s_1, s_2), \gamma_n(s_0, s_1, s_2)\right)\right].$$

We will first prove this for smooth Weierstrass curves, and we will do this by first using the Theorem of the Cube, by considering a *universal point* on a *universal smooth Weierstrass curve*.

1 The universal smooth Weierstrass curve and the proof in the smooth case

Definition 2.2. The *universal smooth Weierstrass curve* $\mathcal{E}$ is the Weierstrass curve over $\operatorname{Spec}\mathcal{R}$ (where $\mathcal{R} = \mathbb{Z}[a_1, a_2, a_3, a_4, a_6, 1/\Delta]$) defined by the homogeneous polynomial

$$\mathcal{W} = y^2z + a_1xyz + a_3yz^2 - x^3 - a_2x^2z - a_4xz^2 - a_6z^3 \in \mathcal{R}[x, y, z].$$

The *universal point* $\mathcal{P}$ on $\mathcal{E}$ is the point in $\mathcal{E}(\mathcal{E})$ corresponding to the identity map on $\mathcal{E}$.

Note here that $\mathcal{P} = \left[\left(\mathcal{O}_{\mathcal{E}}(1), x, y, z\right)\right]$, and that it is universal in the sense that any morphism $S \to \mathcal{E}$ of $\mathcal{R}$-schemes factors through the identity on $\mathcal{E}$. Moreover, $\mathcal{E}$ is universal in the following sense.

Proposition 2.3. *Let E be a smooth Weierstrass curve over S. Then there exist unique morphisms $g\colon E \to \mathcal{E}$ and $h\colon S \to \operatorname{Spec} R$ such the following diagram is commutative, and such that the outer square is Cartesian.*

$$\begin{tikzcd}
E \arrow[r] \arrow[d, "g"'] & \mathbb{P}^2_S \arrow[r] \arrow[d, dashed, "h_{\mathbb{P}^2_{\mathcal{R}}}"'] & S \arrow[d, "h"] \\
\mathcal{E} \arrow[r] & \mathbb{P}^2_{\mathcal{R}} \arrow[r] & \operatorname{Spec}\mathcal{R}
\end{tikzcd}$$

Proof. Mimic the proof of Proposition 1.5. □

We want to consider the multiples of $\mathcal{P}$. Note that for $n \in \mathbb{Z}$, the point $n\mathcal{P}$ is the point corresponding to the multiplication-by-n map μ_n on $\mathcal{E}$. The following is a special case of the Theorem of the Cube.

Theorem 2.4 ([6, Thm. IV.3.3])**.** *Let $n_1, n_2, n_3 \in \mathbb{Z}$. Then there is a canonical isomorphism of $\mathcal{O}_{\mathcal{E}}$-modules*

$$\bigotimes_{I \subseteq \{1,2,3\}} \mu^*_{\sum_{i\in I} n_i} \mathcal{O}_{\mathcal{E}}(1)^{(-1)^{\#I}} = \mathcal{O}_{\mathcal{E}}.$$

Proposition 2.5. *Let $n \in \mathbb{Z}$. Then $\mu_n^*\mathcal{O}_{\mathcal{E}}(1) \cong \mathcal{O}_{\mathcal{E}}(n^2)$.*

Proof. As $\mathcal{E}$ is a smooth Weierstrass curve, we have $\mathcal{O}_{\mathcal{E}}(1) = I^{-3}(0)$; the identity on $\mathcal{E}$ can be given by $\big[\big(\mathcal{O}_{\mathcal{E}}(1), x, y, z\big)\big]$, as well as $\big[\big(I^{-3}(0), X, Y, 1\big)\big]$, where the latter is in the notation of [5, Sect. 2.2]. As μ_{-1} fixes the zero section, we deduce that $\mu_{-1}^{*}\mathcal{O}_{\mathcal{E}}(1) \cong \mathcal{O}_{\mathcal{E}}(1)$. Hence it suffices to show that for all positive $n \in \mathbb{Z}$, we have $\mu_n^{*}\mathcal{O}_{\mathcal{E}}(1) \cong \mathcal{O}_{\mathcal{E}}(n^2)$.

For $n = 2$, we apply Corollary 2.4 with $n_1 = n_2 = 1$, $n_3 = -1$ to see that $\mu_2^{*}\mathcal{O}_{\mathcal{E}}(1) \cong \mathcal{O}_{\mathcal{E}}(4)$. For $n > 2$, our claim follows by induction, by applying Corollary 2.4 with $n_1 = n - 2$, and $n_2 = n_3 = 1$. ☐

As a consequence, we have the following important observation.

Corollary 2.6. *Let $n \in \mathbb{Z}$. Then there exist homogeneous elements α'_n, β'_n, γ'_n in $\mathcal{R}[x,y,z]/(\mathcal{W})$ of degree n^2 such that*

$$\big[\big(\mathcal{O}_{\mathcal{E}}(n^2), \alpha'_n, \beta'_n, \gamma'_n\big)\big] \in \mathcal{E}(\mathcal{E})$$

is the point corresponding to μ_n. These elements are unique up to a common unit in $\mathcal{R}$.

Proof. We simply note that the global sections of $\mathcal{O}_{\mathcal{E}}(n^2)$ are the homogeneous elements of degree n^2 in $\mathcal{R}[x,y,z]/(\mathcal{W})$, and that $\operatorname{Aut}_{\mathcal{O}_{\mathcal{E}}}\big(\mathcal{O}_{\mathcal{E}}(n^2)\big) = \mathcal{O}_{\mathcal{E}}(\mathcal{E})^{\times} = \mathcal{R}^{\times}$. ☐

Translating this observation in terms of the functor of points of $\mathcal{E}$ using the universality of $\mathcal{P}$ then gives the following.

Corollary 2.7. *Let $n \in \mathbb{Z}$. Let T be a scheme over $\mathcal{R}$, and let $P = [(\mathcal{L}, s_0, s_1, s_2)] \in \mathcal{E}(T)$. Then we have*

$$nP = \big[\big(\mathcal{L}^{\otimes n^2}, \alpha'_n(s_0,s_1,s_2), \beta'_n(s_0,s_1,s_2), \gamma'_n(s_0,s_1,s_2)\big)\big].$$

Finally, using the universality of $\mathcal{E}$, we obtain the following.

Corollary 2.8. *Let $n \in \mathbb{Z}$. Let S be a scheme, and let E be a smooth Weierstrass curve over S defined by W as in (1). Let T be an S-scheme, and let $P = [(\mathcal{L}, s_0, s_1, s_2)] \in E(T)$. Then we have*

$$nP = \big[\big(\mathcal{L}^{\otimes n^2}, \alpha'_n(s_0,s_1,s_2), \beta'_n(s_0,s_1,s_2), \gamma'_n(s_0,s_1,s_2)\big)\big].$$

We now finish the proof of Theorem 2.1 in the smooth case.

Proof of Theorem 2.1 (smooth case). We will now show that up to a common unit in $\mathcal{R}$, the triples $(\alpha_n, \beta_n, \gamma_n)$ and $(\alpha'_n, \beta'_n, \gamma'_n)$ are equal. First observe that we have $\beta_0, \beta'_0 \in \mathcal{R}^{\times}$, $\alpha'_0 = 0 = \alpha_0$, and $\gamma'_0 = 0 = \gamma_0$. Hence we may assume that $n \neq 0$.

Consider the smooth Weierstrass curve E over K as defined in Section 1.2. Note that it now suffices to show that $\alpha_n/\alpha'_n = \beta_n/\beta'_n = \gamma_n/\gamma'_n$ as rational functions in $K(E)$, and that this is a unit in $\mathcal{R}$, i.e. of the form $\pm\Delta^i$ for some integer i.

Observe that for all $P = (x : y : 1) \in E(K) - 0$, we have, by Proposition 1.10 and Corollary 2.8,

$$\big(\alpha'_n(P) : \beta'_n(P) : \gamma'_n(P)\big) = nP = \big(\Phi_n(P)\Psi_n(P) : \Omega_n(P) : \Psi_n^3(P)\big)$$

First note that by definition, $\alpha_n/\gamma_n = \Phi_n/\Psi_n^2$ and $\beta_n/\gamma_n = \Omega_n/\Psi_n^3$. Hence $\alpha_n/\gamma_n = \alpha'_n/\gamma'_n$ and $\beta_n/\gamma_n = \beta'_n/\gamma'_n$, as rational functions in $K(E)$. As μ_n is surjective, it follows that $\operatorname{im}\mu_n$ is infinite, hence none of α'_n, β'_n, γ'_n are identically zero. So $\alpha_n/\alpha'_n = \beta_n/\beta'_n = \gamma_n/\gamma'_n$ in $K(E)$. (These are actually rational functions, as both the numerators and denominators are homogeneous of degree n^2.)

Let $\theta_n = \beta_n/\beta'_n$. Then we need to prove that θ_n is in fact a unit in $\mathcal{R}$.

First note that for all $P \in E(K)$, we have $\big(\alpha'_n(P), \beta'_n(P), \gamma'_n(P)\big) \neq (0,0,0)$. Hence θ_n has no poles on E. It follows that θ_n has no zeroes on E either, hence θ_n is constant, i.e. $\theta_n \in K^{\times}$.

As the homogeneous Weierstrass polynomial $\mathcal{W}$ is monic in x, it follows that $\mathcal{R}[x,y,z]/(\mathcal{W})_{n^2}$ is a free $\mathcal{R}$-module, with a basis consisting of monomials. Since θ_n is by definition a quotient of two elements of $\mathcal{R}[x,y,z]/(\mathcal{W})_{n^2}$, which is moreover a constant, it follows that θ_n is in the field of fractions of $\mathcal{R}$.

Now observe that $\mathcal{R}$ is a unique factorisation domain. Hence we can write $\theta_n = f/g$ with $f, g \in \mathcal{R}$ in such a way that f and g have no common factors. Then $f\beta'_n = g\beta_n$ in $\mathcal{R}[x,y,z]/(\mathcal{W})$. Thus g divides all coefficients of β'_n. Now note that for the point $P = (0 : 1 : 0)$, we have $\big(\alpha'_n(P) : \beta'_n(P) : \gamma'_n(P)\big) = (0 : 1 : 0)$, hence the y^{n^2}-coefficient of β'_n is a unit. Therefore, g is a unit as well, which implies that $\theta_n \in \mathcal{R}$.

Finally, note that θ_n divides all coefficients of β_n. But because $\Lambda\Omega_n = 1$ and $\operatorname{ord}_0\Omega_n = -3n^2$, the coefficient of y^{n^2} in β_n must also be 1, as β_n is the homogenisation of a representative of Ω_n. Hence $\theta_n \in \mathcal{R}^\times$, which finishes the proof. $\square$

2 The universal Weierstrass curve and the proof in the general case

As in the case of smooth Weierstrass curves, there exists a *universal Weierstrass curve* $\mathcal{C}$ (as defined in Section 1.1) and a *universal point* $\mathcal{P} = \big[\big(\mathcal{O}_{\mathcal{C}^{\mathrm{sm}}}(1), x, y, z\big)\big] \in \mathcal{C}^{\mathrm{sm}}(\mathcal{C}^{\mathrm{sm}})$, corresponding to the identity map on $\mathcal{C}^{\mathrm{sm}}$. Moreover, the universal smooth Weierstrass curve $\mathcal{E}$ is the complement of the closed subscheme of $\mathcal{C}^{\mathrm{sm}}$ defined by the elliptic discriminant Δ.

Proposition 2.9. *Let $n \in \mathbb{Z}$. Then we have, in $\mathcal{C}^{\mathrm{sm}}(\mathcal{C}^{\mathrm{sm}})$,*

$$n\mathcal{P} = \big[\big(\mathcal{O}_{\mathcal{C}^{\mathrm{sm}}}(n^2), \alpha_n, \beta_n, \gamma_n\big)\big].$$

Proof. Write $n\mathcal{P} = [(\mathcal{L}, s_0, s_1, s_2)] \in \mathcal{C}^{\mathrm{sm}}(\mathcal{C}^{\mathrm{sm}})$. By Proposition 2.5 and Corollary 1.6, we have $\mathcal{L}|_{\mathcal{E}} \cong \mathcal{O}_{\mathcal{C}^{\mathrm{sm}}}(n^2)|_{\mathcal{E}}$. Note that $\mathcal{C}^{\mathrm{sm}}$ is noetherian integral separated regular, therefore the Weil divisor class group is naturally isomorphic to the Picard group.

Note moreover that Δ is irreducible and that $\operatorname{div}\Delta = V(\Delta)$, so $V(\Delta)$ is a principal prime divisor. As $\mathcal{E} = \mathcal{C}^{\mathrm{sm}} - V(\Delta)$, it follows that the natural map $\operatorname{Pic}\mathcal{C}^{\mathrm{sm}} \to \operatorname{Pic}\mathcal{E}$ is an isomorphism. Hence $\mathcal{L} \cong \mathcal{O}_{\mathcal{C}^{\mathrm{sm}}}(n^2)$, so we assume without loss of generality that $\mathcal{L} = \mathcal{O}_{\mathcal{C}^{\mathrm{sm}}}(n^2)$.

Next, we note that by the smooth case of Theorem 2.1, we have

$$u \cdot (s_0|_{\mathcal{E}}, s_1|_{\mathcal{E}}, s_2|_{\mathcal{E}}) = (\alpha_n|_{\mathcal{E}}, \beta_n|_{\mathcal{E}}, \gamma_n|_{\mathcal{E}})$$

for a unique unit u of $\mathcal{R} = \mathcal{A}_\Delta$. As $\mathcal{E}$ is dense in $\mathcal{C}^{\mathrm{sm}}$, we deduce that $u \cdot (s_0, s_1, s_2) = (\alpha_n, \beta_n, \gamma_n)$. Note that the units of $\mathcal{R}$ are $\pm\Delta^i$ with $i \in \mathbb{Z}$. If we pull back both triples via the zero section (i.e. we evaluate all of the sections at $(0 : 1 : 0)$), we see that both $us_0(0,1,0)$ and $us_2(0,1,0)$ are zero, and $us_1(0,1,0) = 1$, as global sections of $\mathcal{O}_{\mathcal{A}}$, i.e. elements of $\mathcal{A}$. Hence we have $s_0(0,1,0) = s_2(0,1,0) = 0$, so as s_0, s_1, s_2 generate $\mathcal{L}$, we deduce that $s_1(0,1,0)$ is a unit in $\mathcal{A}$. So u is a unit in $\mathcal{A}$ as well, i.e. equal to 1 or -1. $\square$

Theorem 2.1 follows from this by the universality of $\mathcal{C}$ and $\mathcal{P}$. Moreover, as the polynomials α_n, β_n, γ_n were defined as homogenisations of $\Phi_n\Psi_n$, Ω_n, Ψ_n^3, it immediately follows that we have the following scheme-theoretic equivalent of Corollary 1.15.

Corollary 2.10. *Let $n \in \mathbb{Z}$. Let S be a scheme, and let C be a Weierstrass curve over S defined by W as in (1). Let T be an S-scheme, and let $P \in C(T)$ be of the form $(x : y : 1)$, such that $a_1y - 3x^2 - 2a_2x - a_4$, $2y + a_1x + a_3$ generate $\mathcal{O}_T$. Then we have*

$$nP = \big(\Phi_n(x,y)\Psi_n(x,y) : \Omega_n(x,y) : \Psi_n^3(x,y)\big).$$

Bibliography

[1] I. Connell. Elliptic curve handbook, February 1999. `http://www.ucm.es/BUCM/mat/doc8354.pdf`.

[2] P. Deligne and M. Rapoport. Les schémas de modules de courbes elliptiques. *Modular Functions of One Variable II, Lecture Notes in Mathematics*, 349:143–316, 1973.

[3] A. Enge. *Elliptic curves and their applications to cryptography — an introduction*. Kluwer, 1999.

[4] A. Grothendieck and J. Dieudonné. Éléments de Géométrie Algébrique IV: Étude locale des schémas et des morphismes de schémas (troisième partie). *Publications mathématiques de l'IHÉS*, 28:5–255, 1966.

[5] N. M. Katz and B. Mazur. *Arithmetic moduli of elliptic curves*, volume 108 of *Ann. Math. Studies*. Princeton University Press, 1985.

[6] M. Raynaud. *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, volume 119 of *Lecture Notes in Mathematics*. Springer-Verlag, 1970.

[7] The Stacks Project Authors. *Stacks Project*. `http://stacks.math.columbia.edu`.